arXiv:chao-dyn/9707019v1 21 Jul 1997

Helicity Transfer in Turbulent Models

L. Biferale[1], D. Pierotti[2] and F. Toschi[3]
[1] *Dipartimento di Fisica, Università di Tor Vergata Via della Ricerca Scientifica 1, I-00133 Roma, Italy and*
Istituto Nazionale di Fisica della Materia, unità di Tor Vergata
[2] *Dipartimento di Fisica, Università dell' Aquila, Via Vetoio 1, I-67010 Coppito, L'Aquila, Italy and*
Istituto Nazionale di Fisica della Materia, unità dell'Aquila
[3] *Dipartimento di Fisica, Università di Pisa, Piazza Torricelli 2, I-56126, Pisa, Italy and*
Istituto Nazionale di Fisica della Materia, unità di Tor Vergata and
Istituto Nazionale di Fisica Nucleare, sezione di Pisa

(February 5, 2008)

Helicity transfer in a shell model of turbulence is investigated. We show that a Reynolds-independent helicity flux is present in the model when the large scale forcing breaks inversion symmetry.
The equivalent in Shell Models of the "2/15 law", obtained from helicity conservation in Navier-Stokes eqs., is derived and tested.
The odd part of helicity flux statistic is found to be dominated by a few very intense events.
In a particular model, we calculate analytically leading and sub-leading contribution to the scaling of triple velocity correlation.

One of the most intriguing problems in three dimensional fully developed turbulence (FDT) is related to the appearance of anomalous scaling laws at high Reynolds numbers, i.e. in the limit when Navier-Stokes dynamics is dominated by the non-linear interactions.

The celebrated 1941 Kolmogorov theory (K41) was able to capture the main phenomenological ideas by performing dimensional analysis based on the energy transfer mechanism. Kolmogorov postulated that the energy cascade should follow a self-similar and homogeneous process entirely dependent on the energy transfer rate, ϵ. This idea, plus the assumption of local isotropy and universality at small scales, led to a precise prediction on the statistical properties of the increments of turbulent velocity fields: $\delta v(l) \sim |v(x+l)-v(x)| \sim (l\epsilon(l))^{1/3}$. From this the scaling of moments of $\delta v(l)$, the structure functions, can be determined in terms of the statistics of $\epsilon(l)$, i.e.

$$S_p(l) \equiv \langle (\delta v(l))^p \rangle = C_p \langle (\epsilon(l))^{p/3} \rangle l^{p/3}, \quad (1)$$

where C_p are constants and the scale l is supposed to be in the inertial range, i.e. much smaller than the integral scale and much larger than the viscous dissipation cutoff. If $S_p(l) \sim l^{\zeta(p)}$ and $\langle \epsilon^p(l) \rangle \sim l^{\tau(p)}$ then

$$\zeta(p) = p/3 + \tau(p/3). \quad (2)$$

In the K41 the $\epsilon(l)$ statistic is assumed to be l-independent, or $\tau(p)=0$, implying $\zeta(p)=\frac{p}{3}, \forall p$. On the other hand, there are many experimental and numerical [1,2] results telling us that K41 scenario for homogeneous and isotropic turbulence is quantitatively wrong. Strong intermittent bursts in the energy transfer have been observed and non trivial $\tau(p)$ set of exponents measured.

Moreover, the problem of investigating scaling properties of observables with the same physical dimensions but with different tensorial structures has been only recently addressed [3,4].

Many different authors have focused their attention on the possible role played by helicity, the second global invariant of 3d Navier-Stokes eqs. [5–8], for determining leading or sub-leading scaling properties of correlation functions in the inertial range.

Recently [4,9], an exact scaling equation for the third order velocity correlations entering in the helicity flux definition has been derived under two hypothesis: (i) there exists a non-vanishing helicity flux, (ii) the flux becomes Reynolds independent in the limit of FDT. This relation predicts a r^2 scaling for a particular third order velocity correlation. The new relation has been called "2/15 law" because of the coefficient appearing in front of the r^2 in analogy with the "4/5 law" derived by Kolmogorov for the third-order structure functions entering in the expression of energy flux. In the "4/5" law the scaling of a different third-order velocity correlation is found to be linear in r.

This simple fact tells us that different velocity correlation with the same physical dimension but with different tensorial structure may have very different scaling properties.

Moreover, even if overwhelming evidences indicate that the main physics is driven by the energy transfer, there can be some sub-leading new intermittent statistics hidden in the helicity flux properties.

Homogeneous and isotropic turbulence has, by definition, always a vanishing mean helical flux. Nevertheless, both fluctuations about the zero-mean in isotropic cases and/or net non-zero fluxes in cases where inversion symmetry is explicitly broken can be of some interest for the understanding of fully developed turbulence.

In this letter, we analyse the helical transfer mechanism in dynamical models of turbulence [10,8,11], built such as to explicitly consider helicity conservation in the inviscid limit.

We give the first strong numerical evidence that a Reynolds-independent helicity flux is present in cases where the forcing mechanism explicitly breaks inversion symmetry.

We confirm that in all cases where two fluxes can coexist in the inertial range, velocity correlations with the same physical dimension but with different transformation properties under inversion symmetry, can show strongly different scaling behaviour.

In the following, we briefly summarize the main motivation behind the introduction of Shell Models for turbulence. We present the equivalent derivation of the "2/15 law" for the helicity flux in our Shell Model language and we conclude with our numerical results about the Reynolds independency of helicity flux and about its statistical intermittent properties.

Shell models have demonstrated to be very useful for the understanding of many properties connected to turbulent flows [12]- [18]. The most popular shell model, the Gledzer-Ohkitani-Yamada (GOY) model ([12]- [18]), has been shown to predict scaling properties for $\zeta(p)$ (for a suitable choice of the free parameters) similar to what is found experimentally.

The GOY model can be seen as a severe truncation of the Navier-Stokes equations: it retains only one complex mode u_n as a representative of all Fourier modes in the shell of wave numbers k between $k_n = k_0 2^n$ and k_{n+1}.

It has been pointed out that GOY model conserves in the inviscid, unforced limit two quadratic quantities. The first quantity is the *energy*, while the second is the equivalent of *helicity* in 3D turbulence [17]. In two recent works [8,11] the GOY model has been generalized in terms of shell variable, u^+_n, u^-_n, transporting positive and negative helicity, respectively. It is easy to realize that only 4 independent classes of models can be derived such as to preserve the same helical structure of Navier-Stokes equations [7]. All these models have at least one inviscid invariants non-positive defined which is similar to the 3D Navier-Stokes helicity. In the following, we will focus on the intermittent properties of a mixture of two of such a models. The mixture is a linear combination of the old GOY model (extended to have u^+, u^-) plus another model which has a different helical interaction and which has already been extensively investigated [11,19]. We focus only on two of the four possible models because they are the only two classes of models which show a clear forward energy cascade (see [11] for more details). The time evolution for positive-helicity shells reads [11]:

$$\dot{u}^+_n = ik_n(A_n[u,u] + xB_n[u,u])^* - \nu k_n^2 u^+_n + \delta_{n,n_0} f^+, \tag{3}$$

with the equivalent eqs, but with all helicity signs reversed, for $\dot{u}^-$. In (3), x defines the relative weights of the two models in the mixture, ν is the molecular viscosity, f^+, f^- are the large scale forcing and $A[u,u]$ and $B[u,u]$ refer to the non-linear terms of the two models. Namely:

$$A_n[u,u] \equiv u^-_{n+2}u^+_{n+1} + b_3 u^-_{n+1}u^+_{n-1} + c_3 u^-_{n-1}u^-_{n-2}, \tag{4}$$
$$B_n[u,u] \equiv u^+_{n+2}u^-_{n+1} + b_1 u^-_{n+1}u^-_{n-1} + c_1 u^-_{n-1}u^+_{n-2}. \tag{5}$$

It is easy to verify that for the choice $b_3 = -5/12, c_3 = -1/24, b_1 = -1/4, c_1 = -1/8$ there are only two global inviscid invariants [11]: the energy, $E = \sum_n(|u^+_n|^2 + |u^-_n|^2)$), and helicity, $H = \sum_n k_n(|u^+_n|^2 - |u^-_n|^2)$.

The equations for the fluxes throughout shell number n are:

$$d/dt \sum_{i=1,n} E_i = k_n < (uuu)^E_n > -\nu \sum_{i=1,n} E_i + E_{in}, \tag{6}$$
$$d/dt \sum_{i=1,n} H_i = k_n^2 < (uuu)^H_n > -\nu \sum_{i=1,n} H_i + H_{in}, \tag{7}$$

where E_i and H_i are the energy and helicity of the nth shell, respectively: $E_i =< |u^+_i|^2 + |u^-_i|^2 >$, $H_i = k_i(< |u^+_i|^2 - |u^-_i|^2 >)$. E_{in} and H_{in} are the input of energy and helicity due to forcing effects, $E_{in} = \Re(< f^+(u^+_1)^* + f^-(u^-_1)^* >)$, $H_{in} = \Re(k_1 < f^+(u^+_1)^* - f^-(u^-_1)^* >)$.
In (6) and (7) we have introduced the triple correlation:

$$\begin{aligned} < (uuu)^E_n > &= (\Delta^+_{n+1} + \Delta^-_{n+1}) + (b_3 + 1/2)(\Delta^+_n + \Delta^-_n) \\ &\quad + x\left((\Gamma^+_{n+1} + \Gamma^-_{n+1}) + (b_1 + 1/2)(\Gamma^+_n + \Gamma^-_n)\right), \quad (8) \\ < (uuu)^H_n > &= (\Delta^+_{n+1} - \Delta^-_{n+1}) + (b_3 + 1/4)(\Delta^+_n - \Delta^-_n) \\ &\quad + x\left((\Gamma^+_{n+1} - \Gamma^-_{n+1}) + (b_1 + 1/4)(\Gamma^+_n - \Gamma^-_n)\right), \quad (9) \end{aligned}$$

and

$$\Delta^+_n = \left\langle \Im(u^+_{n+1}u^-_n u^+_{n-1})\right\rangle, \quad \Gamma^+_n = \left\langle \Im(u^-_{n+2}u^+_{n+1}u^+_n)\right\rangle. \tag{10}$$

Assuming that there exists a stationary state we have $\frac{d}{dt}\Pi^E_n = \frac{d}{dt}\Pi^H_n = 0$, where $\Pi^E_n = k_n\langle(uuu)^E_n\rangle$ and $\Pi^H_n = k_n^2\langle(uuu)^H_n\rangle$. Moreover, in the inertial range we can neglect the viscous contribution in (6) and (7), obtaining:

$$< (uuu)^E_n >= k_n^{-1}E_{in}, \quad < (uuu)^H_n >= k_n^{-2}H_{in}. \tag{11}$$

Supposing that there exist the energy and helicity fluxes, $E_{in} = \epsilon$, $H_{in} = h$ (the latter different from zero only if $f^+ \neq f^-$) and supposing that both are Reynolds-independent, we have in the inertial range:

$$< (uuu)^E_n >\sim k_n^{-1}, \quad < (uuu)^H_n >\sim k_n^{-2}. \tag{12}$$

Relation (11) is the equivalent of what found for helical Navier-Stokes turbulence in [4,9]).

Figure 1 reports results for the helicity flux in a numerical simulations done with two different Reynolds numbers, $Re \sim 10^5$, $Re \sim 10^9$ for a choice of mixture parameter $x = 0.1$ and other numerical inputs as follows, $N = 16$

and 26, $\nu = 10^{-5}$ and $2 \cdot 10^{-9}$. A non-zero helical flux was obtained using a forcing term breaking inversion symmetry: $f^+ = 5 \cdot 10^{-3}(1+i)$, $f^- = f^+/10$. Time marching was obtained by using a slaved Adams-Bashforth algorithm for a number of iterations equal to several thousands of the typical eddy turnover time. A clear inertial range with a non-vanishing helicity flux is detected. The extension of the range where helicity flux is roughly constant scales with the Reynolds number. Moreover, the flux intensity is roughly constant at changing Reynolds number, giving the first evidence that the model can simultaneously support both energy and helicity transfers and that both of them are Reynolds-independent.

Let us remark that this is only possible due to the non-positiveness of helicity; in 2d turbulence, for example, a similar results, concerning enstrophy and energy cascades, is clearly apriori forbidden.

As for the statistics of helicity transfer, we measured the scaling exponents of the moments of energy and helicity fluxes:

$$\Sigma_E^{(p)} \equiv < ((uuu)_n^E)^{(p/3)} > \sim k_n^{-\zeta(p)} \tag{13}$$

$$\Sigma_H^{(p)} \equiv < ((uuu)_n^H)^{(p/3)} > \sim k_n^{-\psi(p)}. \tag{14}$$

As one can see in figure 2 we have that the even part of the two statistics coincides, i.e. $\zeta(2p) = \psi(2p)$. On the other hand, the scaling exponents of odd moments are different.

The difference in odd moments is the signature of strong cancellation effects in the statistics connecting fluctuations at different scales. The picture we have in mind is that the main effect driving turbulent fluctuations is due to the energy cascade process, with its intermittent fluctuations measured by the $\zeta(p)$ exponents. Superimposed to the energy transfer, there are "topological" fluctuations introduced by the asymmetric forcing and measured by the odd-part of the helicity turbulent transfer.

Let us notice that helicity flux fluctuations are much larger then the average helicity flux. This clearly distinguishes the helicity transfer mechanism from the energy transfer mechanism.

In the former, the strong intermittent behaviour shown by odd moments tell us that the odd part of the statistic is dominated by a few very singular structures.

In the case of no-mixture ($x = 0$) one can also exactly calculate sub-leading scaling for the triple correlation $\delta_n^+ \equiv k_n(\Delta_n^+ + \Delta_n^-)$, $\delta_n^- \equiv k_n^2(\Delta_n^+ - \Delta_n^-)$. Indeed, from expressions (8) one obtains after some simple algebra:

$$\delta_n^+ = 2\epsilon \frac{1-y^{n+1}}{1-y}, \quad \delta_n^- = 4h \frac{1-z^{n+1}}{1-z} \tag{15}$$

where $y = -2(b_3 + 1/2)$ and $z = -4(b_3 + 1/4)$. Being both y and z with modulus less then one we recover the asymptotic scaling (12) and one can also control sub-leading correction to it:

$$\Delta_n^+ = k_n^{-1}(\epsilon \frac{1-y^{n+1}}{1-y}) + k_n^{-2}(2h \frac{1-z^{n+1}}{1-z})$$

and

$$\Delta_n^- = k_n^{-1}(\epsilon \frac{1-y^{n+1}}{1-y}) - k_n^{-2}(2h \frac{1-z^{n+1}}{1-z})$$

In conclusion, we have studied a Helical shell model for turbulence with a forcing which explicitly breaks inversion symmetry at large scale.

We have checked that a Reynolds-independent helicity flux establishes in the system, giving the first evidence of very different scaling for triple correlations entering in the energy flux and helicity flux definitions.

The odd part of the helicity flux statistic is found to be strongly intermittent.

For a particular class of models we can also calculate explicitly sub-leading corrections to pure scaling behaviour of typical triple correlation functions.

The existence of sub-leading terms explicitly tell us that scaling laws in turbulent flows must be studied on correlation functions which have a pure projection on the physical relevant quantities.

There are not reasons why similar effects should not be present also in true Navier-Stokes eqs. For example, spurious intermittent corrections can be detected in cases where isotropy is globally or locally violated (as in boundary layers).

Acknowledgements: This work has been partially supported by the INFM (Progetto di Ricerca Avanzata: TURBO).

---

[1] C. Meneveau & K. R. Sreenivasan, J. Fluid Mech. **224**, 429 (1991).

[2] R. Benzi, S. Ciliberto, R. Tripiccione, C. Baudet, C. Massaioli & S. Succi, Phys. Rev. E **48**, R29 (1993).

[3] V. S. L'Vov, E. Podivilov and I. Procaccia, *Invariants for Correlations of Velocity Differences in Turbulent Fields*, preprint in chao-dyn@xyz.lanl.gov ♯ 9706018.

[4] V. S. L'Vov, E. Podivilov and I. Procaccia, *Exact Results for the 3rd Order Correlations of Velocity in Turbulence with Helicity.* preprint in chao-dyn@xyz.lanl.gov ♯ 9705016.

[5] R. H. Kraichnan, J. Fluid Mech. **59**, 745 (1973).

[6] Levich, Physica A **170**, 241 (1991).

[7] F. Waleffe, Phys. Fluids A **4**, 350 (1992).

[8] L. Biferale & R. Kerr, Phys. Rev. E **52**, 6113 (1995).

[9] O. Chkhetiani, JETP Lett. **63**, 808 (1996).

[10] L. Kadanoff, Physics Today, September **11** (1995).

[11] R. Benzi, L. Biferale, R. Kerr and E. Trovatore, Phys. Rev. E **53**, 3541 (1996).

[12] E.B. Gledzer, Sov. Phys. Dokl. **18**, 216 (1973).
[13] M. Yamada & K. Ohkitani, Prog. Theor. Phys. **81**, 329 (1989); M. Yamada & K. Ohkitani, J. Phys. Soc. Jpn. **56**, 4210 (1987); Phys. Rev. Lett **60**, 983 (1988); Prog. Theor. Phys. **79**, 1265 (1988).
[14] M.H. Jensen, G. Paladin and A. Vulpiani; Phys. Rev. A **A43**, 798 (1991).
[15] L. Biferale, A. Lambert, R. Lima and G. Paladin; Physica D **80** 105 (1995).
[16] D. Pisarenko, L. Biferale, D. Courvoisier, U. Frisch and M. Vergassola, Phys. Fluids **A65** 2533 (1993).
[17] L. Kadanoff, D. Lohse, J. Wang and R. Benzi; Phys. Fluids **7** 617 (1995).
[18] E. Leveque and Z.-S. She, Phys. Rev. E **55**, 2789 (1997).
[19] R. Benzi, L. Biferale and E. Trovatore, Phys. Rev. Lett. **77**, 3114 (1996).

FIGURE CAPTIONS

- Figure 1:
  Helicity-flux, $(k_n^2 < (uuu)_n^H >)$, versus k_n for $N = 16$, $\nu = 10^{-5}$ (dashed line) and $N = 26$, $\nu = 2 \cdot 10^{-9}$ (continuos line).

- Figure 2:
  Anomalous exponents for the helicity flux, ψ_p (circles), and for the energy flux, ζ_p (squares), obtained for $N = 26$ and $\nu = 2 \cdot 10^{-9}$.


1.5e-03
5.0e-04
-5.0e-04
-1.5e-03
-2.5e-03
0.0
5.0
10.0
15.0
Π_n^H
$\ln_2(k_n)$


3.0
2.0
1.0
0.0
0.0
2.0
4.0
6.0
8.0
10.0
p
ψ(p)
ζ(p)